Kyle B Potts, M.D., FACS

General, Advanced Laparoscopic & Breast Surgery
Diplomat of the American Board of Surgery

A Professional Corporation
805 W. Acequia, Suite 2A
Visalia, California 93291

Tel: (559) 738-0450 ~ Fax: (559) 738-0460

INSTRUCTIONS FOR PREP WITH CLEAR LIQUID AND ANTIBIOTIC READ CAREFULLY - DO NOT EXCEED RECOMMENDED DOSAGE - SERIOUS SIDE EFFECTS MAY OCCUR

TWO DAYS BEFORE SURGERY

Please read all instructions at least two (2) days before surgery. Your dosage schedule may be adjusted by your physician.

1.	Take Dulcolax as directed.
2.	

If you are taking other medication (including aspirin or aspirin containing products) consult your primary care physicain for additional instructions before beginning this procedure.

DAY BEFORE SURGERY

1. Drink only "clear liquids" for breakfast, lunch and dinner. Solid foods, milk or milk products are not allowed.

"CLEAR LIQUIDS" INCLUDE:

- Juice (apple, white grape)
- Clear broth (beef, chicken or vegetable)
- Tea and coffee (without milk or creamer)
- Water

All of the following that are not colored red or purple:

- Gatorade
- Carbonated and non-carbonated soft drinks
- Kool-Aid or other fruit flavored drinks
- Plain Jell-O without added fruits or toppings
- Ice popsicles
- 2. Take Flagyl 1.0 gram and Neomycin 1.0 gram each at 2 p.m., 3 p.m. and 11 p.m.

DAY OF SURGERY

- No liquids after midnight
- Take your normal medications with a sip of water
- Stop drinking clear liquids 3 hours before the procedure
- Bring your escort/driver with you to your appointment
- · Arrive at your scheduled time

Kyle B Potts, M.D., FACS

General, Advanced Laparoscopic & Breast Surgery
Diplomat of the American Board of Surgery

A Professional Corporation
805 W. Acequia, Suite 2A
Visalia, California 93291
Tel: (559) 738-0450 ~ Fax: (559) 738-0460